

AD_____

GRANT NUMBER DAMD17-97-1-7094

TITLE: Identification and Cloning a Novel Tumor Suppressor Gene
in Breast Cancer

PRINCIPAL INVESTIGATOR: Claudio J. Conti, Ph.D.

CONTRACTING ORGANIZATION: The University of Texas
M.D. Anderson Cancer Center
Houston, Texas 77030

REPORT DATE: December 1999

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

20001013 120

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

| | | | | |
|---|--|---|--|--|
| 1. AGENCY USE ONLY (Leave blank) | | 2. REPORT DATE December 1999 | 3. REPORT TYPE AND DATES COVERED Final (1 Sep 97 - 30 Nov 99) | |
| 4. TITLE AND SUBTITLE Identification and Cloning a Novel Tumor Suppressor Gene in Breast Cancer | | | 5. FUNDING NUMBERS DAMD17-97-1-7094 | |
| 6. AUTHOR(S) Conti, Claudio J., Ph.D. | | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Texas M.D. Anderson Cancer Center Houston, Texas 77030 | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 | | | 10. SPONSORING / MONITORING AGENCY REPORT NUMBER | |
| 11. SUPPLEMENTARY NOTES | | | | |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | 12b. DISTRIBUTION CODE | |
| 13. ABSTRACT (Maximum 200 words) On the basis of an extensive body of cytogenetic literature as well as molecular evidence from our laboratory, we hypothesize that a tumor suppressor gene is located on the long arm of Human Chromosome 7 at the q31.1 band. Furthermore, we propose that inactivation of this tumor suppressor gene plays a role in the development of breast cancer. The long term objectives of the study are to identify and clone the tumor suppressor gene and to determine its function. The objectives for the requested funding period are to provide functional evidence for the existence of this tumor suppressor gene in breast cancer using microcell fusion and to clone the DNA fragment containing the putative tumor suppressor gene using a novel approach based on the introduction of Yeast Artificial Chromosomes (YACs) into breast cancer cells. This novel approach will facilitate the cloning of putative tumor suppressor genes closing the technical gaps that exist between the identification of deleted regions of genome containing putative tumor suppressor genes and the actual cloning of DNA fragments containing the candidate genes. | | | | |
| 14. SUBJECT TERMS Breast Cancer | | | 15. NUMBER OF PAGES 11 | |
| | | | 16. PRICE CODE | |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | 20. LIMITATION OF ABSTRACT Unlimited | |


FOREWORD


Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.


____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


 ☒ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

 ☒ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

 ☒ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.



PI - Signature

11-19-99

Date

Table of Contents

| | Page Number |
|------------------------------------|-------------|
| FRONT COVER | 1 |
| REPORT DOCUMENTATION PAGE..... | 2 |
| FOREWORD | 3 |
| TABLE OF CONTENTS..... | 4 |
| INTRODUCTION (one paragraph)..... | 5 |
| BODY | 6 |
| KEY RESEARCH ACCOMPLISHMENTS | 8 |
| REPORTABLE OUTCOMES | 8 |
| CONCLUSIONS | 9 |
| REFERENCES | 10 |
| BIBLIOGRAPHY | 11 |
| APPENDICES..... | N/A |

INTRODUCTION

Several reports in the literature indicate that karyotypic alterations of chromosome 7 are common in many different types of neoplasia including breast cancer (reviewed in Atkin and Baker, 1991; Pandis, et al. 1993). These alterations include monosomies, trisomies and deletions of different segments of the chromosome. More recently, several laboratories including our own have shown a high frequency of loss of heterozygosity (LOH) in the long arm of chromosome 7 suggesting the existence of a tumor suppressor gene in this region (reviewed in Zenklusen and Conti, 1996). Furthermore, microcell fusion experiments have shown that chromosome 7 can inhibit tumorigenicity or induce senescence in several cell lines (Zenklusen, et al. 1994b; Ogata, et al. 1993). On the basis of this cytogenetic, molecular and functional evidence, we hypothesize that a tumor suppressor gene is located in the long arm of chromosome 7 at the q31.1 band. Furthermore, we propose that inactivation of this tumor suppressor gene plays a role in the development of breast cancer.

The overall objective of these studies is to identify and/or clone this putative tumor suppressor gene using innovative approaches. Thus, we propose to do a number of experiments to develop this approach and to apply them to the cloning of this gene.

The proposed strategy consists of transferring the whole chromosome 7 as well as relevant fragment of this chromosome (as shown by our LOH studies) into breast cancer cell lines. Whole chromosomes can be transferred to the breast cancer cells by microcell fusion as described in previous experiments from our laboratory (Zenklusen, et al. 1994a). To transfer relevant fragments of chromosome 7 we propose to use spheroblast fusion with selected yeast artificial chromosomes (YACs) which contain the markers with higher LOH in breast cancer (Huxley and Gnirke, 1991; Markie, et al. 1993).

We have previously performed microcell fusion with MCF-7 cells obtaining slow growth cultures and abortive colonies. These results are similar to the ones obtained in Dr. Carl Barrett's laboratory (Ogata, et al. 1993) but differ from our previous results in rodent epithelial cells in which introduction of a chromosome 7 suppressed tumorigenicity with minimal effect on the in vitro proliferative potential (Zenklusen, et al. 1994a). Based on these results as well as data from our laboratory indicating that chromosome 7 can inhibit tumorigenicity in prostate cancer cells without affecting growth in tissue culture (Zenklusen, et al. manuscript in preparation), we postulate that two potential tumor suppressor genes may be present in chromosome 7, one affecting proliferation and the other affecting the relationship between the tumor cells and the host. To further explore this possibility we have carried out LOH studies in a number of well characterized breast cell lines using highly polymorphic microsatellite markers to determine the pattern of chromosome alterations in these cell lines and to choose other lines for future studies. We identify two lines that present a series of homozygous markers in the critical region, making them appropriate for these studies.

As we started the microcell fusion experiments, experiments in parallel performed in collaboration with Dr. Jean Claude Zenklusen and Dr. Eric Green (National Human Genome Research Institute, not funded by this grant) identified a very strong candidate gene. This is a new gene, without homology with any other known gene, which is widely expressed in both fetal

and adult tissues. Mutations have been identified in a number of cancer cell lines. Based on these findings, we concentrate our efforts in transfecting a retrofitted bacterial artificial chromosome (BAC) containing this gene (this BAC contains only this gene) and using as a control a BAC from the same region. The BACs have been transfected into two cell lines showing deletion in this region of chromosome 7 and several neo-resistant clones have been obtained from each cell lines and from each BAC. We are in the process of characterizing the clones and expanding to be injected into new mice to check for tumorigenicity. If requested by the DOD, we can submit an additional report in six month. In this report we expect to be able to provide definitive proof that the gene that we have cloned is the tumor suppressor gene in chromosome 7q31.

BODY

A. Experimental Procedures

BAC selection: In order to construct a single fully contiguous series (contig) of BACs, we utilized the YAC based physical map of chromosome 7 previously reported by Bouffard, et al. (1997). The entire region of interest, with boundaries at D7S522 and D7S677, is contained in a single YAC contig. Using the sequence-tagged sites (STS) contained in the YAC contig, a series of BACs were isolated and a minimal tiling path was selected spanning the whole region. In all, nine BACs (RG030h15, RG253b13, RG054j07, RG300c03, RG114a06, RG099b05, RG343p13, GS234b20 and RG068p20) were selected spanning about 1.2 Mb.

BAC retrofitting: The selected BACs were retrofitted to express a mammalian selection marker (*neo*) using the Cre-Lox system according to the protocol by Kim, et al. (1998). Briefly, 500ng of freshly prepared BAC DNA were mixed with 50ng of the retrofitting vector pRETRObac and 1 unit of Cre recombinase (Novagen, Madison, WI) in a total volume of 30 μ l of 1x reaction buffer. After incubation for 60 min. at 37°C, 5 min. at 70°C, 10 min. at RT and 60 min. at 4°C, the total reaction was drop-dialyzed for 3hrs against water on top of Millipore V membranes (13mm, 0.025 μ m pore, #VSWP 01300) floated in a 24 well plates. The whole reaction was then mixed with 20 μ l of GeneHogs electrocompetent cells (Research Genetics, Huntsville, AL) and transferred to a chilled 0.1cm electroporation cuvette. The cells were pulsed at 1.8 kV, 200 ohms, 25 μ F using BioRad Gene Pulser (Hercules, CA). One milliliter of SOC media was then added and the cells were incubated for 60 min. at 37°C, after which aliquots of 200 and 400 μ l were plated in LB plates containing 50 μ g/ml of chloramphenicol and kanamycin, prespread with 50 μ l of 2% X-gal.

The colonies obtained were analyzed for the integrity of the insert by pulse field gel electrophoresis, and correct integration of the retrofitting vector was assessed by PCR as described by Kim, et al. (1998).

Transfection of large DNA fragments (BAC's) Transfections of the BAC's into the breast carcinoma cell lines was performed using the Fugene™ 6 transfection reagent (Boehringer Mannheim). This reagent is a non-liposomal blend which appears to have maximal levels of transfection with reduced toxicity for mammalian cells. Cells were transfected at 70%

confluency following the specification of the manufacturer. The range of plasmid used in these experiments was 1-3 ug.

Tumorigenicity experiments: Tumorigenicity will be evaluated as described in our previous experiments (Zenklusen, et al. 1994b). Briefly, 10⁶ cells will be injected SC in the flank of Balb/c nude mice. Animals are kept until they developed tumors and need to be terminated according with Institutional and Federal guidelines. Tumor mass is estimated with a caliber weekly and recorded for each mouse. Five mice will be injected with each clone.

B. Results

As discussed in the Introduction section, previous experiments had shown that microcell fusion of MCF-7 cells produce abortive colonies. We hypothesize the existence of two suppressor gene is chromosome 7, one homologous to the one postulated by Oshimura and Barret that induce senescence or cell arrest and the other postulated by our laboratory in rodents cells as well as prostate carcinoma and other human tumors, did not alter cell growth in vitro but suppress tumorigenicity in vivo. In the previous report we have also shown that HOMOD analysis (homozygosity mapping of deletions; J. W. Fountain, personal communication) in breast carcinoma cell lines showed two different pattern in chromosome 7 (using contiguous markers of high heterogeneity). Thus, for our experiments we choose two cell lines that have a well defined minimal chromosomal fragment of LOH in our region of interest (D7S496-D7S490 and we have concentrated our efforts into these two cell lines. One of the cell lines, MDA MB231 is an Adenocarcinoma-derived line, and the other is ZR-75-1, a Ductal Carcinoma In Situ-derived line that contain a large area of homozygosity in the critical markers. Both lines have a similar pattern of homozygosity with 8 consecutive markers presenting the same allelic constitution. The probability that the patients were homozygous in this region is 0.000059; therefore we considered that these cell lines present LOH in the critical region and can be utilized for future experiments.

Also, during the last period we have continued our efforts to clone the gene by other conventional methods. For this purpose, we continue our collaboration with Zenklusen and Green (National Human Genome Research Institute). As a result of this collaboration the whole 1.2 Mb contig of the critical region was sequenced and putative genes were identified by Grail, and GeneScan analysis. Matching those predictions with ESTs known to be in the region (by virtue of their location in the GeneMap database or by PowerBlast matching to the dbEST database of the genomic sequence of the BACs) resulted in a complete picture of the genes in the region. Complete cDNAs for the previously unidentified genes were isolated and PCR assays were generated to amplify each exon from a panel of eight prostate/breast carcinoma cell lines for full sequencing. One of these novel genes presented mutations in 4 of the 8 cell lines analyzed. The gene that has been identified has fifteen exons and spans a 130 Kb segment of genomic DNA. The gene is contained in BAC H_RG114a06 and according with our computer prediction and EST content analysis, is the only gene in that BAC. The complete cDNA of this gene, in both directions and each of the two splice variants identified during the cDNA cloning process are being cloned into a mammalian expression vector at the moment. Mutations have been identified in these genes in several cell lines (see Table 1). Also to rule out the possibility

of a tissue culture artifact, we have sequenced 10 samples of colon cancer and we have identified 4 mutations in this set of tumors.

Based on these parallel results we decided to refocus the project in the BAC H_RG114a06 and in the two cell lines that have a bonafide identified deletion in this region. We carry out an experiment in which we transfect BAC H_RG114a06 into the cell lines MDA MB231 and ZR-75-1. As a negative control we transfect BAC H_RG343p13 which is distal to H_RG114a06, of similar size and contains the full Wnt2 gene which is known to be upregulated in breast tumorigenesis, making it a very unlikely TSG candidate.

To perform these experiments we test the sensitivity to G418 (neo) on the cell lines MDA MB231 and ZR-75-1, and determine that optimal concentration for selection are 400 µg/ml for the MDA MB231, and for the ZR-75-1. Under these doses there is no viable cell in the culture after one week of exposure to neo.

Once optimal concentration of neo was determined, we proceed to transfect both cell lines with the selected BAC's. We tested several concentration of DNA and we found that we could get efficient colony formation (approximately 2-4 per plaque), at a concentration of 1µg in a plaque. Colonies were isolated using cloning cylinders and trypsin and expanded sequentially to 24 wells/plate and a six well plate. None of the colonies has presented any apparent impairment in growth and a selected number of colonies have been shown to have the neo markers as detected by PCR. A minimum of 5 colonies has been expanded for each cell line and for each BAC and they will be further characterized using microsatellite markers in that region. To detect tumorigenicity individual colonies will be injected in nude mice. We expect to be able to start injecting clones by the end of the current year and expect to have definitive results of the first clones in March. We expect that the whole experiment will be completed on May or June of 2000.

KEY RESEARCH ACCOMPLISHMENTS:

- By conventional positional cloning we identify a strong candidate gene for the suppressor gene in chromosome 7p31. Mutations were found in breast cancer cell lines as well as in prostate cell lines and samples from colon carcinomas.
- We establish a protocol for efficient transfection of large fragments of DNA into breast cancer cell lines.
- We identify two prototype breast cancer cell lines with deletions in chromosome 7 q31, which are being used to test the capacity of the gene to suppress tumorigenicity.
- We identify a BAC containing the putative tumor suppressor gene and transfect the BAC into two breast carcinoma cell lines. The same cell lines were transfected with an irrelevant BAC to serve as a negative control. This work is in progress. We expect that these experiments will provide proof of concept that tumor suppressor genes can be identified using transfection of large fragments of DNA corresponding to a region of high frequency of LOH.

REPORTABLE OUTCOMES: (List outcomes)

-Manuscripts: A manuscript describing this work is in preparation.

- Patents and licenses applied for/or issued: We have not applied for patents.
- Degrees obtained that are supported by this award: N/A
- Development of cell lines, tissue or serum repositories: Cell lines containing transfected BACs will be made available to the scientific community after the work is published.
- Informatics such as databases and animal models, etc: N/A
- Funding applied for based on work supported by this award: Funding to continue these experiments will be requested through a RO1 mechanism in the next NIH deadline.
- Employment or research opportunities applied for and/or received on experiences/training supported by this award: Two post-doctoral fellows and several technician receive training within this contract.

CONCLUSIONS

This project had two major goals: to develop techniques for functional cloning of tumor suppressor genes and to use this approach to clone a tumor suppressor gene in chromosome 7q31. Both of these goals have been achieved, although in some cases a change of strategy was necessary to adjust the project to technological changes. The first important change was that the whole region of interest was sequenced as part of the Human Genome Project. We collaborate with the group doing the sequencing and computer techniques were used to detect candidate genes. Therefore, the work of cloning was facilitated by this information.

The second strategic change was due to the realization that it was more feasible technically to transfer BAC's rather than YAC's as originally proposed. We have been able to perform transfers of BAC's into mammalian cells in a reproducible manner and we have obtained colonies from two breast cancer cell lines. Cell lines were transfected with a BAC containing the putative tumor suppressor gene and an irrelevant BAC. The tumorigenicity assay of these clones are in progress. We expect that these experiments will provide the proof of concept that this method can be used to clone other tumor suppressor genes or at least narrow down rapidly extensive regions of the genome. Thus, we think that we have made substantial progress to this goal of the project.

As discussed above we have identified a putative tumor suppressor gene. This gene is expressed in most adult tissues and mutations have been found in cell lines and samples of colon adenocarcinomas. We have continued our experiments of BAC transfection using a BAC containing this putative gene. The rationale for continuing the experiments focussed on the BAC H_RG114a06 containing the putative tumor suppressor gene is several fold:

- This particular BAC apparently has the tumor suppressor gene as the only gene. Therefore, if this BAC can suppress tumorigenicity, it will be very solid evidence that this gene is a bona fide tumor suppressor gene in this system because it will be able to suppress tumorigenicity when driven by its own promoter and is embedded in its own genetic background.
- It will accelerate the identification of the tumor suppressor gene because it can be transfected while the complete cDNA and splicing variants of the gene are determined. It also has the advantage that the gene delivered as a BAC will splice in a physiological manner and will have the splicing forms that are normally present in that particular tissue.
- The most important is that as discussed above, it will provide a proof of concept that tumor suppressor genes can be cloned by this functional strategy.

REFERENCES

- Atkin, N.B., Baker, M.C. Numerical chromosome changes in 165 malignant tumors. Evidence for a nonrandom distribution of normal chromosomes. *Cancer Genet. Cytogenet.* 52:113-121, 1991.
- Bouffard, G. G., Idol, J. R., Braden, V. V., Iyer, L. M., Cunningham, A. F., Weintraub, L. A., Touchman, J. W., Mohr-Tidwell, R. M., Peluso, D. C., Fulton, R. S., Ueltzen, M. S., Weissenbach, J., Magness, C. L., Green, E. D. A physical map of human chromosome 7: an integrated YAC contig map with average STS spacing of 79 kb. *Genome Res.*, 7:673-692, 1997.
- Brooks, S.C., Locke, E.R. and Soule, H.D. Estrogen receptor in a human cell line (MCF-7) from breast carcinoma. *J. Biol. Chem.* 248:6251-6253, 1973.
- Huxley, C. and Gnirke, A. Transfer of yeast artificial chromosomes from yeast to mammalian cells. *Bioessays* 13:545-550, 1991.
- Kim, S. Y., Horrigan, S. K., Altenhofen, J. L., Arbieva, Z. H., Hoffman, R., and Westbrook, C. A. Modification of Bacterial Artificial Chromosome Clones Using Cre Recombinase: Introduction of Selectable Markers for expression in eukaryotic cells. *Genome Res.* 8:404-412, 1998.
- Markie, D., Ragousis, J., Senger, G., Rowan, A., Sansom, D., Trowsdale, J., Sheer, D. and Bodmer, W.F. New vector for transfer of yeast artificial chromosomes to mammalian cells. *Somat. Cell Mol. Genet.* 19:161-169, 1993.
- Ogata, T., Ayusawa, D., Namba, M., Takahashi, E., Oshimura, M. and Oishi, M. Chromosome 7 suppresses indefinite division of nontumorigenic immortalized human fibroblast cell lines KMST-6 and SUSM-1. *Mol. Cell Biol.* 13:6036-6043, 1993.
- Oshimura, M., Kugoh, H., Koi, M., Shimizu, M., Yamada, H., Satoh, H. and Barrett, J.C. Transfer of a normal human chromosome 11 suppresses tumorigenicity of some but not all tumor cell lines. *J. Cell Biochem.* 42:135-142, 1990.
- Pandis, N., Heim, S., Bardi, G., Idvall, I., Mandahl, N. and Mitelman, F. Chromosome analysis of 20 breast carcinomas: cytogenetic multiclonality and karyotypic-pathologic correlations. *Genes Chromosom. Cancer* 6:51-7, 1993.
- Zenklusen, J.C., Oshimura, M., Barrett, C.J. and Conti, C.J. Putative tumor suppressor gene in human chromosome 7 inhibits the malignant phenotype of a murine squamous cell carcinoma (SCC) cell line. *Oncogene* 9:2817-2825, 1994a.
- Zenklusen, J.C., Bièche, I., Lidereau, R. and Conti, C.J. CA microsatellite repeat D7S522 is the most commonly deleted region in Human Primary Breast Cancer. *PNAS* 91:12155-12158, 1994b.

Zenklusen, J.C. and Conti, C.J. Cytogenetic, molecular and functional evidence for novel tumor suppressor genes on the long arm of human chromosome 7. Mol. Carcinog. 15:167-175, 1996.

BIBLIOGRAPHY:

Publications and Meeting Abstracts

One manuscript in preparation.

Personnel Receiving Pay From The Research Effort

Cassie Bigbee
Anita Frijhoff
Margaret LaCava
Gabriel Sternik
Sharon Stockman